



Lectures #10 - 11

Second quantization

Evaluation of one and two-particle matrix elements

Coulomb matrix elements: example of two-particle operator

Helium atom: LS coupled states

Chapter 4, pages 95-99, 102-106, Lectures on Atomic Physics

Chapter 11, pages 241-246, Atomic many-body theory

I. Lindgren and J. Morrison


First & second quantization

First quantization


Classical particles are assigned wave amplitudes

Second quantization

Wave fields are “quantized” to describe the problem in terms of “quanta” or particles.



Second quantization: atomic electrons (fermions)





Second quantization: atomic electrons (fermions)

One-electron state $|k\rangle$
Described by the wave function $\Psi_k(q_i)$

Vacuum state
(no electrons)

$$|k\rangle = a_k^\dagger |0\rangle$$

Creation operator



$$\langle 0|0\rangle = 1$$

$$\langle k| = \langle 0| a_k$$

Annihilation operator



$a_k |0\rangle = 0$: “there are no electrons to annihilate in a vacuum”

$$\langle 0| a_k^\dagger = 0$$



Anticommutation relations

$$\{a_i^\dagger, a_j^\dagger\} = a_i^\dagger a_j^\dagger + a_j^\dagger a_i^\dagger = 0$$

$$\{a_i, a_j\} = a_i a_j + a_j a_i = 0$$

$$\{a_i, a_j^\dagger\} = a_i a_j^\dagger + a_j^\dagger a_i = \delta_{ij}$$

Orthonormality of one-electron states:

$$\begin{aligned} a_k |0\rangle &= 0 \\ \langle 0| a_k^\dagger &= 0 \end{aligned}$$

$$\langle i|j\rangle = \langle 0|a_i a_j^\dagger|0\rangle = \langle 0|\delta_{ij} - a_j^\dagger a_i|0\rangle = \delta_{ij}$$

Two-electron states

First quantization:
Slater determinant

Second quantization

$$\Psi_{jk}(q_1, q_2) = \frac{1}{\sqrt{2}} \begin{vmatrix} \psi_j(q_1) & \psi_k(q_1) \\ \psi_j(q_2) & \psi_k(q_2) \end{vmatrix} \quad \Rightarrow \quad |jk\rangle = a_j^\dagger a_k^\dagger |0\rangle$$

Orthonormality of two-electron states:

$$\langle \Psi_{ij}(q_1, q_2) | \Psi_{kl}(q_1, q_2) \rangle = \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}$$

$$\langle ij | kl \rangle = \langle 0 | \underbrace{a_j a_i a_k^\dagger a_l^\dagger}_{\text{reversed order}} | 0 \rangle$$

NOTE: reversed order

Need to evaluate

Orthonormality of two-electron states

$$\langle ij | kl \rangle = \langle 0 | a_j a_i a_k^\dagger a_l^\dagger | 0 \rangle = ?$$

Lets transform $a_j a_i a_k^\dagger a_l^\dagger$ to "normal form":

Move:

Creation operators

Annihilation operators

left

right

Use anticommutation relations to move a_i to the right


$$\begin{aligned} a_j a_i a_k^\dagger a_l^\dagger &= a_j a_l^\dagger \delta_{ik} - a_j a_k^\dagger a_i a_l^\dagger = \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} a_i a_l^\dagger + a_k^\dagger a_j a_i a_l^\dagger \\ &= \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} a_l^\dagger a_i + a_k^\dagger a_j \delta_{il} - a_k^\dagger a_j a_l^\dagger a_i = \\ &= \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} a_l^\dagger a_i + a_k^\dagger a_j \delta_{il} - \delta_{jl} a_k^\dagger a_i + a_k^\dagger a_l^\dagger a_j a_i \end{aligned}$$

Orthonormality of two-electron states

$$\langle ij | kl \rangle = \langle 0 | a_j a_i a_k^\dagger a_l^\dagger | 0 \rangle = ?$$

Lets transform $a_j a_i a_k^\dagger a_l^\dagger$ to "normal order":

Normal
order with
respect to
vacuum



Use anticommutation relations to move a_i and a_j to the right

$$\begin{aligned} a_j a_i a_k^\dagger a_l^\dagger &= a_j a_l^\dagger \delta_{ik} - a_j a_k^\dagger a_i a_l^\dagger = \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} a_i a_l^\dagger + a_k^\dagger a_j a_i a_l^\dagger \\ &= \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} a_l^\dagger a_i + a_k^\dagger a_j \delta_{il} - a_k^\dagger a_j a_l^\dagger a_i = \\ &= \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} a_l^\dagger a_i + a_k^\dagger a_j \delta_{il} - \delta_{jl} a_k^\dagger a_i + a_k^\dagger a_l^\dagger a_j a_i \end{aligned}$$

$$\langle ij | kl \rangle = \langle 0 | a_j a_i a_k^\dagger a_l^\dagger | 0 \rangle$$

Normal order of operators

Why do we want to transform to normal order?

To calculate expectation values: expectation value of normal ordered operators is zero.

$$a_j a_i a_k^\dagger a_l^\dagger = \delta_{ik} \delta_{jl} - a_l^\dagger a_j \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} a_l^\dagger a_i + a_k^\dagger a_j \delta_{il} - \delta_{jl} a_k^\dagger a_i + a_k^\dagger a_l^\dagger a_j a_i$$

$$\begin{aligned} \langle 0 | a_j a_i a_k^\dagger a_l^\dagger | 0 \rangle &= \delta_{ik} \delta_{jl} - \langle 0 | a_l^\dagger a_j | 0 \rangle \delta_{ik} - \delta_{jk} \delta_{il} + \delta_{jk} \langle 0 | a_l^\dagger a_i | 0 \rangle \\ &+ \langle 0 | a_k^\dagger a_j | 0 \rangle \delta_{il} - \delta_{jl} \langle 0 | a_k^\dagger a_i | 0 \rangle + \langle 0 | a_k^\dagger a_l^\dagger a_j a_i | 0 \rangle \\ &= \delta_{ik} \delta_{jl} - \delta_{jk} \delta_{il} \end{aligned}$$

↑ Exactly the result we had in first quantization

Many-particle operators in second quantization

$$a_i^\dagger |0\rangle$$

one-particle state

States

$$a_i^\dagger a_j^\dagger |0\rangle$$

two-particle state

$$a_i^\dagger a_j^\dagger \dots a_n^\dagger |0\rangle$$

N-particle state

} described
by Slater determinants
in first quantization

Operators

First quantization

Second quantization

One-particle operator

$$F = \sum_{i=1}^N f(\mathbf{r}_i) \longrightarrow F = \sum_{i,j} f_{ij} a_i^\dagger a_j$$

Two-particle operator

$$G = \frac{1}{2} \sum_{i \neq j} g(r_{ij}) \longrightarrow G = \frac{1}{2} \sum_{i,j} g_{ijkl} a_i^\dagger a_j^\dagger a_l a_k$$

$$g_{ijkl} = g_{jilk}$$

Many-particle operators in second quantization

First and second quantization description must produce identical results; however, is it more convenient to use second quantization for calculation of properties of many-electron systems as the antisymmetrization properties are carried by the operators rather than the wave functions as in the case of Slater determinants.

$$F = \sum_{i,j} f_{ij} a_i^\dagger a_j$$

$$G = \frac{1}{2} \sum_{i,j} g_{ijkl} a_i^\dagger a_j^\dagger a_l a_k$$

$$f_{ab} = \langle a | f | b \rangle = \int d^3 r \psi_a^\dagger(\mathbf{r}) f(\mathbf{r}) \psi_b(\mathbf{r})$$

$$g_{abcd} = \langle ab | g | cd \rangle = \int d^3 r_1 \int d^3 r_2 \psi_a^\dagger(\mathbf{r}_1) \psi_b^\dagger(\mathbf{r}_2) g(\mathbf{r}_{12}) \psi_c(\mathbf{r}_1) \psi_d(\mathbf{r}_2)$$

The rules for calculating matrix elements are equivalent, for example:

$$\langle ab | G | ab \rangle = \frac{1}{2} \sum_{ijkl} g_{ijkl} \langle 0 | a_b a_a a_i^\dagger a_j^\dagger a_l a_k a_a^\dagger a_b^\dagger | 0 \rangle = g_{abab} - g_{abba}$$

Examples of one and two-particle operators

First quantization

1 $H_0 = \sum_{i=1}^N h_i$

$$h_i = -\frac{1}{2} \nabla_i^2 - \frac{Z}{r_i}$$

2 Coulomb ($1/r_{12}$) interaction: two-particle operator

$$H' = \frac{1}{r_{12}}$$

Second quantization

$$H_0 = \sum_{i=1}^N \epsilon_i a_i^\dagger a_i$$

eigenvalue of h_i

$$H' = \frac{1}{2} \sum_{ijkl} g_{ijkl} a_i^\dagger a_j^\dagger a_l a_k$$

two-electron matrix element
of the Coulomb potential $1/r_{12}$

$$g_{abcd} = \int d^3 r_1 \int d^3 r_2 \psi_a^\dagger(\mathbf{r}_1) \psi_b^\dagger(\mathbf{r}_2) \frac{1}{r_{12}} \psi_c(\mathbf{r}_1) \psi_d(\mathbf{r}_2)$$



He-like systems revisited

$$H_0 = \sum_{i=1}^N \epsilon_i a_i^\dagger a_i$$

Two-electron state $|ab\rangle = a_a^\dagger a_b^\dagger |0\rangle$ is an eigenfunction of H_0 with an eigenvalue $E_{ab}^{(0)} = \epsilon_a + \epsilon_b$

$$H_0 |ab\rangle = (\epsilon_a + \epsilon_b) |ab\rangle$$

However, it is not necessarily an angular momentum eigenstate.

Lets construct eigenstates of $\{H, \mathbf{L}^2, L_z, \mathbf{S}^2, S_z\}$ from $|ab\rangle$ states.

$$\mathbf{L} = \mathbf{l}_a + \mathbf{l}_b$$

$$\mathbf{S} = \mathbf{s}_a + \mathbf{s}_b$$

First, we couple l_a and l_b to construct eigenstates of \mathbf{L}^2, L_z with eigenvalues $L(L+1)$ and M_L

Second, we couple s_a and s_b to construct eigenstates of \mathbf{S}^2, S_z with eigenvalues $S(S+1)$ and M_S

S=0: singlet state
S=1: triplet states

$$s_a = s_b = 1/2 \longrightarrow S = 0, 1$$

He-like systems: construction of coupled states

$$|ab\rangle = a_a^\dagger a_b^\dagger |0\rangle$$

Therefore, the coupled states are given by the sum over magnetic moments which contains two Clebsch-Gordon coefficients:

$$|ab, LM_L, SM_S\rangle = \eta \sum_{m_a m_b} \sum_{\mu_a \mu_b} \left[\begin{array}{c} \downarrow l_a m_a \\ \text{---} \\ \uparrow l_b m_b \end{array} \text{---} \begin{array}{c} \downarrow \frac{1}{2} \mu_a \\ \text{---} \\ \uparrow \frac{1}{2} \mu_b \end{array} \right] a_a^\dagger a_b^\dagger |0\rangle$$

LM_L SM_S

Normalization factor \nearrow

$$\langle ab, LM_L, SM_S | ab, LM_L, SM_S \rangle = \eta^2 \left(1 + (-1)^{L+S} \delta_{n_a n_b} \delta_{l_a l_b} \right)$$

$$\eta = \begin{cases} 1 & n_a \neq n_b, l_a \neq l_b \\ \frac{1}{\sqrt{2}} & \text{Identical orbitals } n_a = n_b, l_a = l_b \end{cases}$$

He-like systems: First-order energy

$$|ab, LM_L, SM_S\rangle = \eta \sum_{m_a m_b} \sum_{\mu_a \mu_b} \left[\begin{array}{c} \downarrow l_a m_a \\ \text{---} LM_L \text{---} \\ \uparrow l_b m_b \end{array} \right] \left[\begin{array}{c} \downarrow \frac{1}{2} \mu_a \\ \text{---} SM_S \text{---} \\ \uparrow \frac{1}{2} \mu_b \end{array} \right] a_a^\dagger a_b^\dagger |0\rangle$$

The first-order energy is given by

$$E^{(1)} = \langle ab, LM_L, SM_S | H' | ab, LM_L, SM_S \rangle$$

$$H' = \frac{1}{2} \sum_{ijkl} g_{ijkl} a_i^\dagger a_j^\dagger a_l a_k$$

$$g_{abcd} = \int d^3 r_1 \int d^3 r_2 \psi_a^\dagger(\mathbf{r}_1) \psi_b^\dagger(\mathbf{r}_2) \frac{1}{r_{12}} \psi_c(\mathbf{r}_1) \psi_d(\mathbf{r}_2)$$



Coulomb matrix element

First, let's consider Coulomb matrix element

$$g_{abcd} = \int d^3 r_1 \int d^3 r_2 \psi_a^\dagger(\mathbf{r}_1) \psi_b^\dagger(\mathbf{r}_2) \frac{1}{r_{12}} \psi_c(\mathbf{r}_1) \psi_d(\mathbf{r}_2)$$

The functions ψ are given by $\psi_{nlm}(\mathbf{r}) = R_{nl}(r) Y_{lm}(\theta, \phi)$


The $1/r_{12}$ can be expanded as

$$\frac{1}{r_{12}} = \sum_{k=0}^{\infty} \frac{4\pi}{2k+1} \frac{r_{<}^k}{r_{>}^{k+1}} \sum_{q=-k}^k Y_{kq}^*(\theta_1, \phi_1) Y_{kq}(\theta_2, \phi_2)$$

This expression may be re-written using C-tensors defined by

$$C_q^k(\hat{r}) = \sqrt{\frac{4\pi}{(2k+1)}} Y_{kq}(\theta, \phi)$$

$$\frac{1}{r_{12}} = \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} = \sum_{k=0}^{\infty} \frac{r_{<}^k}{r_{>}^{k+1}} \sum_{q=-k}^k (-1)^q C_q^k(\hat{r}_1) C_{-q}^k(\hat{r}_2)$$



Coulomb matrix element

We now substitute the expressions for ψ and $1/r_{12}$ back into our matrix element and separate dr and $d\Omega$ integrals

$$\frac{1}{r_{12}} = \sum_{k=0}^{\infty} \frac{r_{<}^k}{r_{>}^{k+1}} \sum_{q=-k}^k (-1)^q C_q^k(\hat{r}_1) C_{-q}^k(\hat{r}_2)$$

$$\psi_{nlm}(\mathbf{r}) = R_{nl}(r) Y_{lm}(\theta, \phi)$$

$$g_{abcd} = \int d^3 r_1 \int d^3 r_2 \psi_a^\dagger(\mathbf{r}_1) \psi_b^\dagger(\mathbf{r}_2) \frac{1}{r_{12}} \psi_c(\mathbf{r}_1) \psi_d(\mathbf{r}_2)$$

$$R_k(abcd)$$

$$= \sum_{k=0}^{\infty} \int_0^{\infty} dr_1 r_1^2 \int_0^{\infty} dr_2 r_2^2 R_{n_a l_a}(r_1) R_{n_b l_b}(r_2) \frac{r_{<}^k}{r_{>}^{k+1}} R_{n_c l_c}(r_1) R_{n_d l_d}(r_2)$$

radial integral

$$\times \sum_{q=-k}^k (-1)^q$$

$$\int d\Omega_1 Y_{l_a m_a}(\theta_1, \phi_1) C_q^k(\theta_1, \phi_1) Y_{l_c m_c}(\theta_1, \phi_1)$$

$$\langle l_a m_a | C_q^k | l_c m_c \rangle$$

$$\int d\Omega_2 Y_{l_b m_b}(\theta_2, \phi_2) C_{-q}^k(\theta_2, \phi_2) Y_{l_d m_d}(\theta_2, \phi_2)$$

$$\langle l_b m_b | C_{-q}^k | l_d m_d \rangle$$

Coulomb matrix element

$$g_{abcd} = \sum_{k=0}^{\infty} R_k(abcd) \sum_{q=-k}^k (-1)^q \langle l_a m_a | C_q^k | l_c m_c \rangle \langle l_b m_b | C_{-q}^k | l_d m_d \rangle$$

Next, we use Wigner-Eckart theorem for both of the matrix elements:

$$g_{abcd} = \sum_{k=0}^{\infty} R_k(abcd) \sum_{q=-k}^k (-1)^q \begin{array}{c} \uparrow l_a m_a \\ | \\ k \ q \\ | \\ l_c m_c \end{array} \begin{array}{c} \uparrow l_b m_b \\ | \\ k \ -q \\ | \\ l_d m_d \end{array} \langle l_a || C^k || l_c \rangle \langle l_b || C^k || l_d \rangle$$


We use $\begin{array}{c} \uparrow l_b m_b \\ | \\ k \ -q \\ | \\ l_d m_d \end{array} = (-1)^{k-q} \begin{array}{c} \uparrow l_b m_b \\ | \\ k \ q \\ | \\ l_d m_d \end{array}$

Note: k and q are integers

Coulomb matrix element

$$\begin{aligned}
 g_{abcd} &= \sum_{k=0}^{\infty} R_k(abcd) \sum_{q=-k}^k (-1)^k \begin{array}{c} \uparrow l_a m_a \\ | \\ \text{---} k q \\ | \\ \downarrow l_c m_c \end{array} \begin{array}{c} \uparrow l_b m_b \\ | \\ \text{---} k q \\ | \\ \downarrow l_d m_d \end{array} \langle l_a \| C^k \| l_c \rangle \langle l_b \| C^k \| l_d \rangle \\
 &= \sum_{k=0}^{\infty} \boxed{(-1)^k R_k(abcd) \langle l_a \| C^k \| l_c \rangle \langle l_b \| C^k \| l_d \rangle} \begin{array}{c} \uparrow l_a m_a \\ | \\ \text{---} k \\ | \\ \downarrow l_c m_c \end{array} \begin{array}{c} \uparrow l_b m_b \\ | \\ \text{---} \\ | \\ \downarrow l_d m_d \end{array} + \\
 &\quad \downarrow \\
 &\quad \boxed{X_k(abcd)} \\
 &\quad \downarrow \\
 &\quad \boxed{g_{abcd} = \sum_{k=0}^{\infty} \begin{array}{c} \uparrow l_a m_a \\ | \\ \text{---} k \\ | \\ \downarrow l_c m_c \end{array} \begin{array}{c} \uparrow l_b m_b \\ | \\ \text{---} \\ | \\ \downarrow l_d m_d \end{array} + X_k(abcd)}
 \end{aligned}$$

Summary: Coulomb matrix element (non-relativistic case)



$$g_{abcd} = \sum_{k=0}^{\infty} - \text{Diagram} + X_k(abcd)$$

$$X_k(abcd) = (-1)^k R_k(abcd) \langle l_a \| C^k \| l_c \rangle \langle l_b \| C^k \| l_d \rangle$$

$$R_k(abcd) = \int_0^{\infty} dr_1 r_1^2 \int_0^{\infty} dr_2 r_2^2 R_{n_a l_a}(r_1) R_{n_b l_b}(r_2) \frac{r_{<}^k}{r_{>}^{k+1}} R_{n_c l_c}(r_1) R_{n_d l_d}(r_2)$$

$$\langle l_1 \| C^k \| l_2 \rangle = (-1)^{l_1} \sqrt{(2l_1 + 1)(2l_2 + 1)} \begin{pmatrix} l_1 & k & l_2 \\ 0 & 0 & 0 \end{pmatrix}$$

Back to He-like systems

$$|ab, LM_L, SM_S\rangle = \eta \sum_{m_a m_b} \sum_{\mu_a \mu_b} \left[\begin{array}{c} \downarrow l_a m_a \\ \text{---} LM_L \text{---} \\ \uparrow l_b m_b \end{array} \right] \left[\begin{array}{c} \downarrow 1/2 \mu_a \\ \text{---} SM_S \text{---} \\ \uparrow 1/2 \mu_b \end{array} \right] a_a^\dagger a_b^\dagger |0\rangle$$

Assume
 $l_a = l_{a'}$
 $l_b = l_{b'}$

$$E^{(1)} = \langle ab, LM_L, SM_S | H' | ab, LM_L, SM_S \rangle$$

$$= \eta^2 \sum_{m_a m_b} \sum_{\mu_a \mu_b} \sum_{m_{a'} m_{b'}} \sum_{\mu_{a'} \mu_{b'}} \left[\begin{array}{c} \downarrow l_a m_a \\ \text{---} LM_L \text{---} \\ \uparrow l_b m_b \end{array} \right] \left[\begin{array}{c} \downarrow 1/2 \mu_a \\ \text{---} SM_S \text{---} \\ \uparrow 1/2 \mu_b \end{array} \right] \left[\begin{array}{c} \downarrow l_a m_{a'} \\ \text{---} LM_L \text{---} \\ \uparrow l_b m_{b'} \end{array} \right] \left[\begin{array}{c} \downarrow 1/2 \mu_{a'} \\ \text{---} SM_S \text{---} \\ \uparrow 1/2 \mu_{b'} \end{array} \right]$$

$$\times \left(g_{a'b'ab} \delta_{\mu_a \mu_{a'}} \delta_{\mu_b \mu_{b'}} - g_{a'b'ba} \delta_{\mu_a \mu_{b'}} \delta_{\mu_b \mu_{a'}} \right)$$

$$H' = \frac{1}{2} \sum_{ijkl} g_{ijkl} a_i^\dagger a_j^\dagger a_l a_k; \quad \langle ab | G | ab \rangle = g_{abab} - g_{abba}$$

Sum over μ 's: Term 1

Term 1: $\sum_{\mu_a \mu_b} \sum_{\mu_a' \mu_b'} \delta_{\mu_a \mu_a'} \delta_{\mu_b \mu_b'}$

$= (2S + 1) \text{ (circular diagram)} = 1$

We used orthogonality relation to evaluate this diagram

$$\text{(circular diagram)} = \frac{1}{2j_3 + 1} \delta_{j_3' j_3} \delta_{m_3' m_3}$$

Sum over μ 's: Term 2

$$\text{Term 2: } \sum_{\mu_a \mu_b} \sum_{\mu_a' \mu_b'} \delta_{\mu_a \mu_b'} \delta_{\mu_b \mu_a'} - \begin{array}{c} \downarrow \frac{1}{2} \mu_a \\ \text{---} SM_S \\ \uparrow \frac{1}{2} \mu_b \end{array} - \begin{array}{c} \downarrow \frac{1}{2} \mu_{a'} \rightarrow \mu_b \\ \text{---} SM_S \\ \uparrow \frac{1}{2} \mu_{b'} \rightarrow \mu_a \end{array}$$

$$= (2S + 1)(-1)^{S+1} \begin{array}{c} \frac{1}{2} \\ \text{---} SM_S \text{---} \bigcirc \text{---} SM_S \\ \text{+} \quad \quad \quad \text{-} \\ \frac{1}{2} \end{array} = -(-1)^S$$

Extra phase factor \uparrow

First-order energy: He-like systems

$$E^{(1)} = \eta^2 \sum_{m_a' m_b'} \sum_{m_a m_b} \sum_k - \text{diagram} - \text{diagram}$$

Assume

$$l_a = l_a'$$

$$l_b = l_b'$$


$$\times \left(- \text{diagram} + X_k(abab) + (-1)^S - \text{diagram} + X_k(abba) \right)$$

where we substituted Coulomb matrix elements g into our formula


$$g_{abcd} = \sum_{k=0}^{\infty} - \text{diagram} + X_k(abcd)$$

Sum over m's: Term 1

$$\sum_{m_a' m_b'} \sum_{m_a m_b}$$



rotate
to the right



- 1) Remove triangle (get 6-j symbol)
- 2) Use orthogonality relation
(get factor $1/(2L+1)$)

$$= (-1)^{L+k+l_a+l_b} \begin{Bmatrix} l_a & l_b & L \\ l_b & l_a & k \end{Bmatrix}$$

Sum over m's: Term 2

$$\sum_{m_a' m_b'} \sum_{m_a m_b} - \begin{array}{c} \downarrow l_a m_a \\ \text{---} LM_L \\ \downarrow l_b m_b \end{array} - \begin{array}{c} \downarrow l_a m_{a'} \\ \text{---} LM_L \\ \downarrow l_b m_{b'} \end{array} - \begin{array}{c} \uparrow l_a m_{a'} \\ \text{---} k \\ \uparrow l_b m_b \end{array} + \begin{array}{c} \uparrow l_b m_{b'} \\ \text{---} l_a m_a \end{array} =$$

$$= (-1)^{k+l_a+l_b} \begin{Bmatrix} l_a & l_b & L \\ l_a & l_b & k \end{Bmatrix}$$

The calculation is very similar to Term1, only need to switch a and b lines in the first angular diagram.

Final expression:

First order energy for He-like system

$$E^{(1)} = \eta^2 \sum_k \left\{ (-1)^{L+k+l_a+l_b} \begin{Bmatrix} l_a & l_b & L \\ l_b & l_a & k \end{Bmatrix} X_k(abab) \right. \\ \left. + (-1)^{S+k+l_a+l_b} \begin{Bmatrix} l_a & l_b & L \\ l_a & l_b & k \end{Bmatrix} X_k(abba) \right\}$$

$$X_k(abcd) = (-1)^k R_k(abcd) \langle l_a \| C^k \| l_c \rangle \langle l_b \| C^k \| l_d \rangle$$

$$R_k(abcd) = \int_0^\infty dr_1 r_1^2 \int_0^\infty dr_2 r_2^2 R_{n_a l_a}(r_1) R_{n_b l_b}(r_2) \frac{r_{<}^k}{r_{>}^{k+1}} R_{n_c l_c}(r_1) R_{n_d l_d}(r_2)$$

$$\langle l_1 \| C^k \| l_2 \rangle = (-1)^{l_1} \sqrt{(2l_1+1)(2l_2+1)} \begin{pmatrix} l_1 & k & l_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{Bmatrix} = 1$$

Special case: if $l_a = l_b = 0$ (s states) $\rightarrow \langle s \| C^k \| s \rangle = \delta_{k0}$